


Name: Key

Physics Quiz 1

1. 2.0 moles of a monatomic ideal gas initially are at state **A**, with a pressure of $P_1 = 3.0 \times 10^5$ Pa and volume of $V_1 = 6.0 \times 10^{-2}$ m³. The gas can be taken directly to state **B** by the constant-temperature path **A**→**B** depicted, or stepwise through state **C**, by path **A**→**C** followed by path **C**→**B**. m³ m³



- a. The path **A**→**B** is carried out at constant temperature. The word for a constant temperature process is Isothermal.
- b. The path **A**→**C** is carried out at constant volume. The word for that is isochoric.
- c. The path **C**→**B** is carried out at constant pressure. The word for that is isobaric.
- d. What is the temperature of the gas at state **A**?

$$\text{Ideal gas } PV = nRT, \text{ so } T = PV/(nR) = \frac{(3 \times 10^5 \text{ J/m}^3)(6 \times 10^{-2} \text{ m}^3)}{(2 \text{ mol})(8.314 \text{ J/(mol K)})} = 1082.5 \text{ K}$$

This seemed high to me, so I made the online version 4 mol gas, making the temperature half as much, or 541.3 K.

- e. What is the internal energy of the gas at state **A**?

$$\text{Monatomic ideal gas } U = 3/2 nRT = 3/2 PV = 3/2(3 \times 10^5 \text{ J/m}^3)(6 \times 10^{-2} \text{ m}^3) \\ = 27 \times 10^3 \text{ J} = 27 \text{ kJ}$$

- f. The volume of the gas at state **B** is $V_2 = 9.0 \times 10^{-2}$ m³. What is the pressure P_2 at state **B**?

$$PV = nRT \text{ and } T \text{ is unchanged, so } P_1 V_1 = P_2 V_2 \text{ thus } P_2 = P_1 V_1 / V_2 \\ = (3 \times 10^5 \text{ J/m}^3)(6 \times 10^{-2} \text{ m}^3) / (9 \times 10^{-2} \text{ m}^3) = 2 \times 10^5 \text{ J/m}^3 = 2 \times 10^5 \text{ Pa}$$

- g. What is the internal energy of the gas at state **B**?

Same temperature as state **A**, so same $U = 27$ kJ.

- h. What is the internal energy *change* of the gas along path **A**→**B**?

Zero.

- i. What is the sign (+, -, 0) of the work done by the gas along path **A→B**?
The gas expands, so it does work on the surroundings. (+)
- j. What is the work done by the gas along path **A→B**?
Work done by an ideal gas in an isothermal step:
 $W = nRT \ln(V_2/V_1) = (18 \text{ kJ}) \ln(9/6) = (18 \text{ kJ})(0.4055) = 7298 \text{ J}$
- k. What is the sign (+, -, 0) of the heat absorbed by the gas along path **A→B**?
Here, $Q = W$ (Why?) so (+)
- l. What is the heat absorbed by the gas along path **A→B**?
 $Q = W$, so $Q = 7298 \text{ J}$.
- m. What is the sign (+, -, 0) of the work done by the gas along path **A→C**?
0
- n. What is the work done by the gas along path **A→C**?
0
- o. What is the temperature of the gas at state **C**?
 $PV = nRT$ and here V is constant so $V = V_1 = nRT_A/P_1 = nRT_C/P_2$. Then
 $T_C = T_A P_2/P_1 = (1082.5 \text{ K})(2/3) = 701.7 \text{ K}$.
If there are 4 moles of gas, the temperature is half this,
360.9 K.
- p. What is the internal energy of the gas at state **C**?
 $U = 3/2 nRT = 3/2 P_2 V_1 = 3/2 (12 \text{ kJ}) = 18 \text{ kJ}$
- q. What is the internal energy *change* of the gas along path **A→C**?
 $\Delta U = U_C - U_A = 18 \text{ kJ} - 27 \text{ kJ} = -9 \text{ kJ}$
- r. What is the sign (+, -, 0) of the heat absorbed by the gas along path **A→C**?
Zero work, decrease in U , so negative Q . (-)

- s. What is the heat absorbed by the gas along path **A**→**C**?

$$\Delta U = Q - W, \text{ so } Q = \Delta U + W = -9 \text{ kJ} + 0 = -9 \text{ kJ}.$$

- t. What is the sign (+, -, 0) of the work done by the gas along path **C**→**B**?

Expansion, so the system does work. (+)

- u. What is the work done by the gas along path **C**→**B**?

$$\text{For an isobaric process, } W = P\Delta V = (2 \times 10^5 \text{ J/m}^3)(3 \times 10^{-2} \text{ m}^3) = +6 \text{ kJ}$$

- v. What is the change in the internal energy of the gas along path **C**→**B**?

$$\Delta U = U_B - U_C = 27 \text{ kJ} - 18 \text{ kJ} = +9 \text{ kJ}$$

- w. What is the sign (+, -, 0) of the heat absorbed by the gas along path **C**→**B**?

+

- x. What is the heat absorbed by the gas along path **C**→**B**?

$$\Delta U = Q - W, \text{ so } Q = \Delta U + W = 9 \text{ kJ} + 6 \text{ kJ} = 15 \text{ kJ}$$

2. The First law of thermodynamics can be written $\Delta U = Q - W$.

- a. What is U in this formula? What does it mean?

Internal energy, the total molecular kinetic and intermolecular potential energy.

- b. What is Q in this formula? What does it mean?

Heat absorbed by the system.

- c. What is W in this formula? What does it mean?

Work done by the system.